



UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU  
GRAMIN KRISHI MAUSAM SEWA(GKMS)  
AMFU OF IMD, BENGALURU



**AGROMET-ADVISORY BULLETIN**

Date: **14.10.2022**

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

**The forecast is valid for Bengaluru Rural district**  
**Weather forecast (Valid from 15-10-2022 to 19-10-2022)**

**Forecast summary:**

Parameters	15.10.2022	16.10.2022	17.10.2022	18.10.2022	19.10.2022
Rainfall (mm)	44	25	25	16	13
Max Temp Trend ( °C)	28	28	27	27	27
Min Temp Trend ( °C)	20	20	20	20	20
Total cloud cover (octa)	7	7	7	7	7
Relative humidity (%)Max	96	96	96	98	98
Relative humidity (%)Min	90	90	90	92	92
Wind speed(Km/hr)	2	3	3	2	3
Wind Direction (Degrees)	135	101	117	115	288

**Moderate to high rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 27.0-28.0°C and Minimum of 20.0°C. Relative humidity 96-98 % during morning hrs and 90-92 % during noon is expected. Wind speed is 2-3 km/hr.**

**Weather Based Agro Advisories**

**Crop information and Crop Stages of the major Kharif/Rabi crops**

District	Kharif crops				Horticulture crops	
Bangalore Rural (BR)	Groundnut	Redgram	Finger millet	Maize	Grape	Mango
	PF,PM	V,F	V,T,E	Ts, CI,GF,FD	-	--

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Transplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting, IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching, CI: Cob Initiation

**Agromet Advisory:**

Crop/ Component	Stage/ Condition	Pest and Disease	Agro advisories
General			1. Chitta rainstar starts from October 11 <sup>th</sup> and remains up to October 23 <sup>rd</sup> . The normal rainfall of Chitta rainstar is 55.4 mm.
Kharif crops	Tillering stage/Ear head formation stage/flowering/pod formation and maturity stage		The following crops are suggested for sowing. Finger millet Ragi (Indaf-5, 7,9, ML-365, KMR-301), Horse gram- PHG-9, KBH-1, Niger, Field bean-HA-3 and 4 and sunflower-KBSH-1,41,42, 44 & 53, cowpea (KBC-1, TVX-944 and PKB-4 for vegetable purpose). 1. Suggested for intercultivation for Ragi crop 2. To control of leaf folder insect in Redgram to spray Methomyl 40SP, 2 gram/litre of water before flowering of crop. 3. If Wilt diseases noticed in Redgram field –Drenching Carbendazim 50 WP 2 g/litre of water. 4. Remove and burn the infected Wilt diseases Redgram plant in

		field.
<b>Horticulture crop</b>		
<b>Mango</b>		1. Paclobutrazol soil application @ 5 g a.i per plant is recommended for uniform flowering behavior in Mango crop. Paclobutrazol is effective not only in flower induction but also in early and off season flower induction which thereby maintains regularity and synchronization in flowering.
<b>Sericulture</b>		
		1. Maintain the optimum room temperature in Sericulture unit.
<b>Dairy</b>		<p>➤ Congenial climate and abundant fodder availability facilitate improved Dairy, Piggery, Poultry, Sheep and Goat rearing.</p> <p><b>Feeding of lactating cow</b></p> <p>➤ Proper feeding of dairy cattle should envisage minimum wastage of nutrients and maximum returns in respect of milk produced.</p> <p>➤ A concentrate mixture made up of protein supplements such as oil cakes, energy sources such as cereal grains (Maize, Jowar), tapioca chips and laxative feeds such as brans (rice bran, wheat bran, gram husk) is generally used.</p> <p>➤ Mineral mixture containing major and all the trace elements should be included at a level of 2 percent.</p> <p>1. Dietary fiber for milking cow should be 17 % and NDF (Neutral Detergent Fiber ) 22 %</p>

**AMFU of IMD  
Bengaluru**

**Important Note:** Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: [mausam.imd.gov.in](http://mausam.imd.gov.in)