



UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU  
GRAMIN KRISHI MAUSAM SEWA(GKMS)  
AMFU OF IMD, BENGALURU



**AGROMET-ADVISORY BULLETIN**

Date: **03.10.2022**

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

**Significant past weather for the preceding week**

Parameter	29.9.2022	30.9.2022	01.10.2022	02.10.2022	03.10.2022
Rainfall (mm)	0	0.2	26.4	6.0	0
Max. temp(°C)	27.4	26.8	25.0	24.8	26.2
Min.Temp(°C)	19.6	17.6	17.0	16.4	16.6
Sky condition(Octas)	4	4	8	6	4
Relative humidity(%) 0830 hours	95	93	98	94	87
Relative humidity(%) 1730 hours	61	73	68	60	...
Wind Speed (kmph)	3.7	4.2	4.4	3.6	4.5
Wind Direction	270	320	230	230	230

**Weather forecast (Valid from 04-10-2022 to 08-10-2022)**

**Forecast summary:**

Parameters	04.10.2022	05.10.2022	06.10.2022	07.10.2022	08.10.2022
Rainfall (mm)	1	0	1	2	5
Max Temp Trend ( °C)	27	29	29	29	29
Min Temp Trend ( °C)	20	20	20	20	20
Total cloud cover (octa)	7	7	7	8	8
Relative humidity (%)Max	87	87	87	89	89
Relative humidity (%)Min	69	69	69	71	71
Wind speed(Km/hr)	6	5	5	6	6
Wind Direction (Degrees)	324	304	270	246	238

**Light rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 27.0-29.0°C and Minimum of 20.0°C. Relative humidity 87-89 % during morning hrs and 69-71 % during noon is expected. Wind speed is 5-6 km/hr.**

**Weather Based Agro Advisories**

**Crop information and Crop Stages of the major Kharif/Rabi crops**

District	Kharif crops				Horticulture crops	
	Groundnut	Redgram	Finger millet	Maize	Grape	Mango
Bangalore Urban (BU)	PF	V	EV,V,T	Ts, Cl,GF	-	--

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Tranplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tilling., Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting  
IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching

**Agromet Advisory:**

Crop/ Component	Stage/ Condition	Pest and Disease	Agro advisories
General			1. Hasta rainstar starts from September 27 <sup>th</sup> and remains up to

		October 10 <sup>th</sup> . The normal rainfall of Hasta rainstar is 105.8 mm.
<b>Kharif crops</b>	Vegetative stage And Tillering stage	<p>The following crops are suggested for sowing.</p> <p>Finger millet Ragi (Indaf-5, 7,9, ML-365, KMR-301), Horse gram-PHG-9, KBH-1 , Niger, Field bean-HA-3 and 4 and sunflower-KBSH-1,41,42, 44 &amp; 53, cowpea (KBC-1, TVX-944 and PKB-4 for vegetable purpose).</p> <ol style="list-style-type: none"> <li>1. Suggested for intercultivation for Ragi crop</li> <li>2. Wherever over population of crop plants is existing, undertake thinning operation for better availability of scarce soil moisture to the reduced population of crop plants.</li> <li>3. To control of leaf folder insect in Redgram to spray Methomyl 40SP , 2 gram/litre of water before flowering of crop.</li> <li>4. If Wilt diseases noticed in Redgram field –Drenching Carbendazim 50 WP 2 g/litre of water.</li> <li>5. Remove and burn the infected Wilt diseases Redgram plant in field.</li> </ol>
<b>Horticulture crop</b>		
<b>Mango</b>		<ol style="list-style-type: none"> <li>1. Paclobutrazol soil application @ 5 g a.i per plant is recommended for uniform flowering behavior in Mango crop. Paclobutrazol is effective not only in flower induction but also in early and off season flower induction which thereby maintains regularity and synchronization in flowering.</li> </ol>
<b>Sericulture</b>		
		<ol style="list-style-type: none"> <li>1. Maintain the optimum room temperature in Sericulture unit.</li> </ol>
<b>Dairy</b>		<ul style="list-style-type: none"> <li>➤ Congenial climate and abundant fodder availability facilitate improved Dairy, Piggery, Poultry, Sheep and Goat rearing.</li> </ul> <p><b>Feeding of lactating cow</b></p> <ul style="list-style-type: none"> <li>➤ Proper feeding of dairy cattle should envisage minimum wastage of nutrients and maximum returns in respect of milk produced.</li> <li>➤ A concentrate mixture made up of protein supplements such as oil cakes, energy sources such as cereal grains (Maize, Jowar), tapioca chips and laxative feeds such as brans (rice bran, wheat bran, gram husk) is generally used.</li> <li>➤ Mineral mixture containing major and all the trace elements should be included at a level of 2 percent.</li> </ul> <ol style="list-style-type: none"> <li>1. Dietary fiber for milking cow should be 17 % and NDF (Neutral Detergent Fiber ) 22 %</li> </ol>

**AMFU of IMD  
Bengaluru**

**Important Note:** Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: [mausam.imd.gov.in](http://mausam.imd.gov.in)