

**UNIVERSITY OF AGRICULTURAL SCIENCES, BENGALURU &  
INDIAN METEOROLOGICAL DEPARTMENT**



**GRAMIN KRISHI MAUSAM SEWA  
AMFU, OFRS, NAGANAHALLI,  
MYSURU - 570003**



Date: 12-01-2024

**AGRO-ADVISORY BULLETIN FOR KODAGU DISTRICT**

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

**Weather forecast for the next five days (From 13-01-2024 to 17-01-2024)**

Parameter	13.01.2024	14.01.2024	15.01.2024	16.01.2024	17.01.2024
Rainfall (mm)	0	0	0	0	0
Max. Temp. (°C)	27	27.5	28.6	28.6	30.1
Min. Temp. (°C)	17.5	16.5	16.7	16.5	17
Sky condition (Octas)	3	3	3	3	3
Relative humidity (%) 0830 hours	99	99	99	99	99
Relative humidity (%) 1730 hours	80	80	80	80	80
Wind Speed (kmph)	10	7	6	5	5
Wind Direction	72	73	75	117	195

**Forecast Summary**

As forecast received from IMD, Cloudy sky with no rainfall may be expected from 13.01.2024 to 17.01.2024 in Kodagu district. The day temperature is expected to be 27-30.1°C & night temperature is expected 16.5-17.5°C. The relative humidity in the morning hours is expected to be 99% & afternoon relative humidity is expected to be in the range of 80%. Wind speed expected to be 5-10 km/ hr.

**Recommendations to the farmers:**

Crop	Pest/Disease	Damage symptoms	Control measures
<ul style="list-style-type: none"> <li>No rainfall expected for next 3-5 days. Farmers are advised to go for harvesting of Agriculture Crops which attained the maturity.</li> <li>The grains of the harvested crops should be properly dried by retaining moisture percentage as mentioned below for long storage &amp; also minimize the store pest damage Cereals 11-12 %, Pulses-9%, Oil seeds-8% and Vegetable seeds 5-6%.</li> <li>Farmers are advised to take up the sowing of pulses like black gram, green gram and also the green manure crop sun hemp to utilize residue moisture after harvest of the paddy in Paddy field.</li> <li>In paddy fields where there is no residue moisture, in such fields farmers can take up deep</li> </ul>			

**ploughing, so that the soil pathogens will die and in turn it will help to reduce pest incidence for the next paddy crop and which will facilitate conservation of rainwater.**

<b>Coconut</b>	<b>Rugose whitefly</b>	<b>The immature and adult whiteflies have a sucking feeding habit, feeding on the under surfaces of the leaflets. Extensive feeding of the insect leads to the excretion of honey dew which encourages growth of the fungus Capnodium sp. which affects the photosynthetic efficiency of the plant. Sooty mould (Capnodium sp.) growth on the leaf surface can be flaked out by spraying 2.5% of maida paste solution mixed with detergent/ Khadi soap @ 5g/l. In severe cases, spray only neem oil 0.5% or NSKE 5% and avoid spraying any form of insecticides.</b>
<b>Animals, Sericulture and Poultry</b>	<b>Due to low temperature in morning and night hours, Maintain the optimum room temperature in Sericulture, Poultry and Dairy unit by providing electric bulb for creating warm room temperature.</b>	

- Download “**DAMINI**” app to get early warning on lightening and take precautions based on the alert given by the application.
- Kindly download “**MAUSAM**” APP for location specific forecast & warning & “**MEGHDOOT**” APP for Agromet advisory
- This information is available in the website: [mausam.imd.gov.in](http://mausam.imd.gov.in)

For any information farmers can contact **Dr.C.Ramachandra**, Senior Farm Superintendent/ **Dr. Sumanth Kumar.G.V**, Research Associate over phone No. 0821-259126/ 9535345814.

**AMFU of IMD,  
Naganahalli, Mysuru**