



UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU
GRAMIN KRISHI MAUSAM SEWA(GKMS)
AMFU OF IMD, BENGALURU



AGROMET-ADVISORY BULLETIN

Date: 13.10.2023

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

The forecast is valid for Bengaluru urban district.

Significant past weather for the preceding week

Parameter	06.10.2023	07.10.2023	08.10.2023	09.10.2023	10.10.2023
Rainfall (mm)	0	0	0	22.8	42.2
Max. temp(°C)	29.8	30.2	29.6	30.2	29.0
Min.Temp(°C)	20.0	20.4	19.8	20.4	18.0
Sky condition(Octas)	2	4	4	2	4
Relative humidity(%) 0830 hours	91	90	83	83	81
Relative humidity(%) 1730 hours	58	55	56	62	--
Wind Speed (kmph)	2.5	2.3	2.5	1.7	2.7
Wind Direction	230	230	230	320	230

Weather forecast (Valid from 14-10-2023 to 18-10-2023)

Forecast summary:

Parameters	14.10.2023	15.10.2023	16.10.2023	17.10.2023	18.10.2023
Rainfall (mm)	0	0	4	4	4
Max Temp Trend (°C)	29	29	29	29	30
Min Temp Trend (°C)	20	20	20	20	21
Total cloud cover (octa)	5	5	5	5	5
Relative humidity (%)Max	70	70	72	72	70
Relative humidity (%)Min	63	63	65	65	63
Wind (Km/hr)	3	2	2	3	4
Wind Direction (Degrees)	90	103	105	101	90

Light rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 29.0-30.0°C and Minimum of 20.0-21.0°C. Relative humidity 70-72 % during morning hrs and 63-65 % during noon is expected. Wind speed is 2-4 km/hr.

Weather Based Agro Advisories

Crop information and Crop Stages of the major Kharif/Rabi crops

District	Kharif crops				Horticulture crops	
	Groundnut	Redgram	Finger millet	Maize	Grape	Mango
Bangalore Urban (BU)	PI,PF	F,PF	EV,V,E	M,H	-	-

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Transplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting, IBI: Inflorescence Bud initiation, PP(V): Pod Picking Vegetable, F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching

Agromet Advisory:

Crop/Component	Stage/Condition	Pest and Disease	Agro advisories
General			<ul style="list-style-type: none"> Chitta rainstar starts from October 11th and remains up to October 23rd. The normal rainfall of Chitta rainstar is 55.4 mm. Deep ploughing is recommended for conserve the soil moisture in field. Intercultural operation recommended for conserving soil moisture

		<p>and removal of weeds.</p> <ul style="list-style-type: none"> • Suggested for intercultivation for Ragi crop • Mulching is recommended to conserve the soil moisture in standing crops.
Kharif crops	Flowering stage and Maturity stage	<p>The following crops are suggested for sowing.</p> <p>Finger millet Ragi (Indaf-5, 7,9, ML-365, KMR-301), Horse gram-PHG-9, KBH-1 , Niger, Field bean-HA-3 and 4 and sunflower-KBSH-1,41,42, 44 & 53, cowpea (KBC-1, TVX-944 and PKB-4 for vegetable purpose).</p> <ul style="list-style-type: none"> • Earthing up makes better availability of soil moisture to the crop rows through conversion of the land into ridges and furrow system. • Wherever over population of crop plants is existing, undertake thinning operation for better availability of scarce soil moisture to the reduce population of crop plants. • To control leaf folder insect in Redgram spray Methomyl 40SP , 2 gram/litre of water before flowering of crop. • Wilt disease noticed: Redgram field-Drench with Carbendazim 50 WP 2 g/litre of water. • For Biological control of pod borer insect in Redgram 200 LE/acre HANPV 400 litre with teepol (1 ml/lit.). Pheromone traps for <i>Helicoverpa armigera</i> 12/ha • Remove and burn the infected Wilt diseased Redgram plant in field.
Mango		<ul style="list-style-type: none"> • Deep ploughing is recommended for conserve the soil moisture and removal of weeds in field. • Paclobutrazol soil application @ 5 g a.i per plant is recommended for uniform flowering behavior in Mango crop. Paclobutrazol is effective not only in flower induction but also in early and off season flower induction which thereby maintains regularity and synchronization in flowering.
Animal Husbandry		
Dairy		<p>Feeding of lactating cow</p> <ul style="list-style-type: none"> • Proper feeding of dairy cattle should envisage minimum wastage of nutrients and maximum returns in respect of milk produced. • A concentrate mixture made up of protein supplements such as oil cakes, energy sources such as cereal grains (Maize, Jowar), tapioca chips and laxative feeds such as brans (rice bran, wheat bran, gram husk) is generally used. • Mineral mixture containing major and all the trace elements should be included at a level of 2 percent. • Dietary fiber for milking cow should be 17 % and NDF (Neutral Detergent Fiber) 22 %

**AMFU of IMD
Bengaluru**

Important Note: Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: mausam.imd.gov.in