



UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU
GRAMIN KRISHI MAUSAM SEWA(GKMS)
AMFU OF IMD, BENGALURU



AGROMET-ADVISORY BULLETIN

Date: **24.05.2022**

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

Significant past weather for the preceding week

Parameter	20.5.2022	21.5.2022	22.5.2022	23.5.2022	24.5.2022
Rainfall (mm)	1.8	0.8	0	0	0
Max. temp(°C)	22.6	24.8	26.4	28.2	29.4
Min.Temp(°C)	16.0	16.4	19.2	19.6	19.8
Sky condition(Octas)	8	6	6	4	3
Relative humidity(%) 0830 hours	91	86	84	88	86
Relative humidity(%) 1730 hours	77	65	55	54	--
Wind Speed (kmph)	8.0	7.3	7.3	8.8	5.8
Wind Direction	140	230	230	270	230

The forecast is valid for Bengaluru Urban district

Weather forecast (Valid from 25-05-2022 to 29-05-2022)

Forecast summary:

Parameters	25.5.2022	26.5.2022	27.5.2022	28.5.2022	29.5.2022
Rainfall (mm)	1	2	0	0	0
Max Temp Trend (°C)	32	32	30	32	32
Min Temp Trend (°C)	21	21	20	20	21
Total cloud cover (octa)	6	7	7	4	4
Relative humidity (%)Max	82	82	87	87	85
Relative humidity (%)Min	55	55	58	57	55
Wind speed(Km/hr)	3	4	4	3	3
Wind Direction (Degrees)	257	259	248	249	249

Light rain forecasted by IMD, Bangalore during next 5 days. The Maximum temperature ranges from 30.0-32.0°C and Minimum of 20.0-21.0°C. Relative 82-87 % during morning hrs and 55-58 % during noon is expected. Wind speed is 3-4 km/hr.

Weather Based Agro Advisories

Crop information and Crop Stages of the major Kharif/Rabi crops

District	<i>Kharif crops</i>				<i>Horticulture crops</i>	
	Groundnut	Redgram	Finger millet	Maize	Grape	Mango
Bangalore Urban (BU)	--	--	--	--	-	FD,M

G: Germination, S: Sowing, EV: Early vegetative, VG: Vegetative growth, TR: Tranplanting, PI: Peg initiation, FLI: Flag leaf initiation, F: Flowering, PF: Pod formation, PM: Pod Maturity, T: Tillering,, Ts: Taselling, E: Ear head emergence, GF: Grain filling, H: Harvesting
IBI: Inflorescence Bud initiation , PP(V): Pod Picking Vegetable , F& FS: Flowering to fruit setting, FD: Fruit Development, H: Harvesting, M: Maturation, B: Branching

Agromet Advisory:

Crop/ Component	Stage/ Condition	Pest and Disease	Agro advisories
General			<ul style="list-style-type: none"> Conditions are favorable for further advance of Southwest Monsoon over some parts of Southwest Arabian Sea, some more parts of

		<p>Southeast Arabian Sea, Maldives & Comorin area, South & East central Bay of Bengal and some parts of Northeast Bay of Bengal during next 48 hours.</p> <ul style="list-style-type: none"> • Rohini rainstar starts from May 25th and remains upto June 7th. The normal rainfall of Rohini rainstar is 61.9 mm. • The farmers are procure good quality of seeds and fertilizers. Take up the germination test for Kharif sowing. • Take up land preparation and strengthen the bund with fodder grass which will facilitate conservation of rainwater due to pre-monsoon showers. Ploughing across the slope will make insitu soil moisture conservation. • Taken up summer plough to expose the soil to kill the different stages of insect and weeds. • Time for application tank silt to increase soil fertility. • Sufficient rainfall received in May month sowing of long duration crops like Redgram BRG-1, 2, 4 &5, TTB-7 and Castor DCH-9, 32 &177 is recommended.
Horticulture crop		
Mango	Fruit Development and Maturity stage	<ol style="list-style-type: none"> 1. In mango, fruit fly damage is also noticed. suitable control measure by spraying Carbaryl 4 g/litre of water 2. Harvesting of mango without dropping ground level using mango net. 3. Mango growers should harvest matured fruit and try to leave one or two inches of stem in the mango bunch, which helps to avoid sap burn disease. Afterward, the mango fruit should be wrapped by using papers and then keep it in plastic crates. After within 24 hours of harvesting, the crates are placed on top of each other with 5 to 6 create half-feet apart and then take 300 ml. of water in small plastic bottles, add 10 ml of ethylene to water, then after adding 10 g sodium hydroxide (caustic soda). Then seal the tarpaulin cover tightly without aeration for 48 to 60 hours, then tarpaulin should be removed and left out for aeration. Ethylene should be kept in 8 to 10 glasses for to ripen 1 ton of fruit. Natural ripening will help to get good quality fruit and better market price.
Sericulture		
	1. Maintain the optimum room temperature in Sericulture unit.	

**AMFU of IMD
Bengaluru**

Important Note: Farmers are informed to use the APPs & Videos related to Weather information: MEGHDOOT, MAUSAM AND DAMINI APPS. This information is available in the website: mausam.imd.gov.in